

Investigation of the effect of POMC peptides on dermal fibroblast migration and its role in wound healing

SamiaTayeb Hawisa¹, Hana Fahelbun², Salah Bahroun³and Naziha Mansuri³

¹Department of Pathology and Immunology, Dental Faculty, Tripoli University; ²Department of Histology, Medical Faculty, Zawia University; ³National Medical Research Center, Zawia

Running title:**Enhanced fibroblast migration by POMC peptides**

Key words: **Fibroblasts, immunohistochemistry, wound healing, POMC peptides**

Corresponding author: SamiaTayebHawisa e-mail: samiahaw@hotmail.com

Abstract:

The human skin contains several different fibroblast subtypes including dermal sheath (DS), dermal papilla (DP) and inter-follicular dermis cells (DF). It has been shown that quicker healing and less scar formation is related to proportionally fewer numbers of DS and DP fibroblasts. TGF- β plays a crucial role in wound healing by attracting fibroblasts and promoting their synthesis of collagens I, III, and V, proteoglycans, fibronectin and other extracellular matrix (ECM) components; and that the POMC peptides α -MSH/ ACTH may ameliorate hypertrophic scarring by antagonizing TGF- β . The aim of this project was to assess the role of POMC peptides α -MSH/ ACTH in modulating migratory activity of fibroblasts *in vitro* using scratch assay with or without POMC peptides and the transwell assay in the presence of the extracellular protein fibronectin. Transwell assay showed that fibronectin increased fibroblast sub-types migration with DS cells migrated through fibronectin transwells more effectively than other fibroblast sub-types. POMC peptides, ACTH induced the greatest increase in migration with fibronectin coated wells. In addition, scratch assay showed that the fibroblast migration was significant in the presence of the POMC peptides.

Introduction

Skin is complex, multilayered organ, which include appendages (hair follicles, sebaceous glands, sweat glands, apocrine glands) and consists of mixed cell types and extracellular components and contains three layers: epidermis, dermis and hypodermis (Freinkel and Woodley, 2001). The epidermis is highly organized kinetic structure in which cells divide, move and differentiate and contains many cell types including keratinocytes, melanocytes,

Langerhans cells and Merkel cells. The dermis is the connective tissue component of the skin, which is much thicker than the epidermis and composed primarily of fibroblasts, collagen, elastin, and adipocytes (Slominski, et al, 2000a&b; MacNeil 2007). It protects the body from mechanical injury and provides its elasticity and tensile strength. There is great interest in

fibroblast regulation because improved activity is important for wound healing and during formation of hypertrophic scars (Freinkel and Woodley, 2001). The dermis and epidermis interact together to maintaining the properties of both tissues and in repairing the skin in the case of trauma. In human beings there are clear differences in wound healing responses between hairy and non-hairy body sites (Jahoda, et al, 2001a). The contribution of the hair follicle-derived dermal cells results in qualitatively enhanced dermal repair (Jahoda et al, 2001b). Normal wound healing involves complex and dynamic events leading to the repair of injured tissues including rapid proliferation of fibroblasts, followed by collagen biosynthesis (Montagna, et al, 1964). Human skin contains several different fibroblast subtypes including interfollicular dermal fibroblasts (DF), the follicular dermal sheath fibroblasts (DS), and the follicular dermal papilla fibroblasts (DP). The dermal sheath cells maintain and regenerate the dermal papilla, a key for hair growth; in addition these cells may have other roles in the wound healing and repair of skin dermis after injury (Jahoda, et al, 2001b). Fibroblasts degrade the provisional matrix via metalloprotease (MMPs) and respond to cytokines by proliferating and synthesizing new extracellular matrix (ECM) to restore the wounded tissue with a connective tissue

scar. Transforming growth factor TGF- β contributes to the scar development by attracting fibroblasts and promoting their synthesis of collagens I, III, and V, proteoglycans, fibronectin and other ECM components. TGF- β concurrently inhibits proteases while enhancing protease inhibitors, favouring matrix accumulation (Wahl, 1999; Branton and Kopp, 1999). The fibroblasts are stimulated to migrate into the wound area, proliferate and produce the matrix proteins fibronectin, hyaluronan (HA) and later collagen and proteoglycans. Once within the wound environment, fibroblasts proliferate and establish to create the new ECM, which is important for the repair progression and supports more ingrowth of cells (Grinnell, 1994). Fibroblasts differentiate into myofibroblasts which help the wound begins to contract (Eichler, et al, 2006; Romo, et al, 2006), contraction can last for several weeks (Mulvaney, et al, 1994) and continues even after the wound is completely re-epithelialized. Myofibroblasts are stimulated by growth factors and ECM fibronectin to move along it linked to fibrin in the provisional ECM in order to reach the wound edges (Hinz, 2006). The proopiomelanocortin (POMC) peptides ACTH, Alpha-melanocyte stimulating hormone (α -MSH), and β -MSH and β -endorphin peptides are detected in normal and

pathological cells as melanocytes, keratinocytes, Langerhans cells, and mononuclear dermal inflammatory cells by immunocytochemistry. The human dermal endothelial cells and fibroblasts produce and released α -MSH and ACTH immunoreactivity into the medium (Slominski, et al, 2000a; Millington, 2006). α -MSH is the most potent melanotropic peptide arising from the proteolytic cleavage of POMC, and is produced in the skin. Moreover, the melanocortin receptor 1 (MC-1R), which is bound by α -MSH and ACTH, is expressed on some types of cells in the skin as keratinocytes, dendritic cells, macrophages and endothelial cells (Luger, et al, 2000). α -MSH peptides and its receptors regulate melanogenesis but can also affect non-pigmentary processes, such as inflammation, apoptosis and sebogenesis (Ruta, et al. 2007). They have effect on the regulation of immune and inflammatory responses, hair growth, exocrine gland activity and extracellular matrix composition (Luger, et al, 1995). Cytoprotective function for this hormone or peptides in defence of the skin from

Methods and materials:

Fibroblast cell culture and maintenance

Fibroblast cells were maintained in growth medium, containing 500 ml RPMI 1640, 50ml of 10% Foetal calf serum, 5ml of Penicillin/streptomycin and 5ml of L-

exogenous factors or stress e.g. would occur following UV exposure or exposure to agents inducing inflammation or oxidative stress (Eves, et al, 2000). Production of melanocortins and expression of their receptors suggest a complex autocrine and/or paracrine regulatory system whose disorder always affects skin homeostasis. These effects of the melanocortin are mediated by melanocortin-1 receptors which are most ubiquitously expressed by human skin cells (Böhmand Luger, 2000). Alpha-MSH is a basic tridecapeptide, the amino acid sequence of α -MSH is identical to the 1-13 (N-terminal) amino acid sequence of ACTH although α -MSH has slight straight control on glucocorticoid secreted from the adrenal gland (Catania, et al. 1993). α -MSH originate primarily in the pituitary, but it is also secreted in low concentrations in the central nervous system, the skin, and other sites. This peptide was named for its effect on pigmentation in amphibian skin; although it is important to other functions due to its tissues distribution in higher organisms (Catania, et al. 1993).

glutamine at an appropriate temperature and gas mixture (5% CO₂, 95% Air at 37⁰C), in cell incubator in humid atmosphere.

Immunocytochemistry:

For immunohistochemistry fibroblast cells (5×10^3 cells/well) were allowed to attach overnight in the incubator then washed with PBS, drained and incubated in ice-cold methanol for 10 min at -20°C . The cells were then fixed in 4% paraformaldehyde in PBS for 10 min at room temperature (this was used for the melanocortin 1 receptor as this receptor is located on the cell surface). The cells were drained and washed by PBS 200 μl /well, 3 times for 5 minutes. Following incubation with 200 μl of 10% Goat serum (GS) for 90 min to block any non-specific sites cells were washed three times in 200 μl PBS for 5 min then incubated with 100 μl of diluted primary antibody (Table 1) per chamber overnight at 4°C . After washing in PBS, 100 μl of pre diluted secondary antibody per chamber was added, and incubated for 15 min. The cells washed with

200 μl PBS for 5 min, three times and drained. 100 μl of streptavidin peroxidase solution was added to the cells and incubated for 20 minutes after washing three times with 200 μl PBS for 5 min, 200 μl of distilled water was added and incubated for 2 min then 200 μl of chromogen was added in each well and incubated for further 10 min (the negative control was used to check for non-specific binding and to determine duration exposure) which result in a red precipitate at the antigen site. During the incubation period the progress of the reaction was monitored under bright field of light microscope. The chambers were washed with distilled water 200 μl per chamber, to stop the chromogen reaction. After mounting with pre warmed glycerol and applying cover slip, pictures were taking of the cells visualize the type and location of any staining.

Table 1: primary antibodies plating of cells onto 8-well glass chamber slides

Antibodies	Concentration
Fibroblast marker	1 : 200 (positive control) 1 : 500 1 : 1000
Melanocortin-1 receptor	1 : 10
Vimentin	1 : 100 1 : 400

Fibroblasts Migration test: The “Scratch Assay”

For scratch assay cells were seeded in 24-well plates (Nunc, Langensfeld, Germany) at 3×10^4 cells/well, at 80% confluence, a scratch

wound using a sterile pipette tip was performed and the size of the gap was measured; cells were incubated with 500 μl of working dilution POMC peptides

(α -MSH, ACTH), each at a concentration of 10^{-7} M and 10^{-8} M, were added to the cells, 500 μ l of full media and basal media were added to the cells as positive and negative controls respectively and the closing of the scratch wound was measured for 72 hours or

until wound was closed. The cells were then stained with (Toluidin blue 1% in Borax) at room temperature, to facilitate digital photography of the wound edges under low magnification using a colour digital camera (Nikon Coolpix) connected with the light microscope.

Trans-well Migration of the Fibroblasts in presence or absence of Fibronectin:

To assess cell migration fibroblasts were placed in the top chamber of transwell migration chambers (8 or 5 μ m; BD Biosciences, San Jose, CA). The lower chamber was filled with 500 μ l of either conditioned medium or control medium with or without 50 μ l containing 5μ g/cm² of fibronectin. The fibronectin solution was aspirated from the transwell plates on the second day. Then the cell suspension was prepared and 8×10^3 cells/well was seeded gently into the transwells

after the counting. 150 μ l of cell suspension was added to each transwell, and 500 μ l of the conditioned media (α -MSH, ACTH) as a chemo-attractant, the full media, and the basal media, were added to the bottom of each well. Cells that had not migrated to the lower chamber were removed from the upper surface of the transwell membrane with a cotton swab. Migrating cells on the lower membrane surface were fixed, stained using Methanol/acetone containing 0.1% crystal violet and Hoechst 33258 dye, and photographed.

Results

Transwell Assay:

In transwell assay the subpopulation of the fibroblast (dermal papilla, dermal sheath, interfollicular cells types) from the same individual (male 46 years old) were assessed and used at density of 8×10^3 cells per well. The chemoattractant factor fibronectin secreted by the fibroblast cells

affects the cells migration, and was investigated in this project. It was shown that fibronectin was a potent chemotactic signal for migration. Figure 1 showed increased the migration rate of DS fibroblast cells with the fibronectin compared with those without fibronectin.





Figure 1: DS fibroblast migration through the membrane coated by fibronectin (a) or uncoated by fibronectin (b) in full media (serum supplement) X2 objective magnifications.

a. with fibronectin b. without fibronectin

Effects of the POMC peptides on fibroblasts migration:

It was also investigated how fibroblast migration is affected by the addition of POMC peptides (α -MSH / ACTH) as a chemotactic stimulus for fibroblasts,

results showed that there increase of migration of cells grown on wells containing α -MSH/ ACTH peptides (Figure 2 and 3).



Figure 2: DS fibroblast migration through the membrane coated by fibronectin in basal media (serum starved) containing α -MSH 10^{-7} M. X2 objective magnifications.

a. α -MSH 10^{-7} M with fibronectin

b. α -MSH 10^{-7} M without fibronectin




Figure 3: DS fibroblast migration through the membrane coated by fibronectin in basal media (serum starved) containing ACTH 10^{-7} M peptide in the presence of fibronectin. X2 objective magnifications

a. ACTH 10^{-7} M with fibronectin

b. ACTH 10^{-7} M without fibronectin

Scratch Assay:

The rate of wound closure and cellular migration of the fibroblast cells was important because if follicular fibroblasts are involved in the wound healing it would be useful to discover their potential rate of migration into a zone of injury. The average distance between leading cell edges photography

of mechanical scratched distance of approximately 2.0 mm, at t0, then after 72 hours from the scratch (t72), and then after 144 hours from the scratch (t144).



Figure 4: Typical central score created on the floor of 24- well plate onto a monolayer of fibroblast, at seeding density of 3×10^4 /well. The photographs were taking after the mechanical scratch directly at (t0), by digital camera (A camera 1/15, F3.2), x 2 objective magnification.




Figure 5: Fibroblast migration at (t72) and at (t 144) with full media (serum supplement), x4 magnification

a. at (t72)

b. at (t 144)

The rate and extent of cell movement was greater with the basal media containing POMC peptides than with basal media alone. Moreover, the migration of fibroblasts was greater in the present of ACTH peptide than in the present of α -MSH peptide (Figure 6).





Figure 6: Fibroblast migration at (t 144). (a) Basal media only. (b) Basal media with α -MSH.10-7 M, (c) basal media with ACTH 10-7 M. X4 objective magnifications.

a. Basal media only

b. Basal media with α -MSH.10-7 M



c. basal media with ACTH 10-7 M

Immunocytochemistry

It was important to know about the immunocytochemistry of the fibroblast cells *in vitro*, to find out the localization of proteins in the three fibroblast subtypes cells, within chemically fixed cells, primary antibodies that identify proteins were added and incubated with the cells, and detected by fluorescence or colorimetric process and visualized by microscopy. The immunocytochemistry was done for the interfollicular dermal fibroblast (DF) cells for female 54 years, and also for fully matched cell types (DF/ DS/ DP) of the male 46 years, but for many reasons there was insufficient number of these cells, as the cells become contaminated at the beginning of the cell

culture and then the human fibroblast cells enter stage of arrested (no more growth) so the female 54 was used.

In contrast with the positive control (fibroblast marker antibody 1:200) the fibroblast cells were tested positive for the protein vimentin (positive staining) with the two concentrations (1:500, 1:1000) it is giving the red colour after adding the chromogen.

The cells also exhibit the melanocortin-1 Receptor (MC1R) expression at a concentration of (1:10) only, but did not reveal any staining of (1:25/1:50/1:100). The results are shown in figure x 10 objective magnification.



Figure 7: Immunocytochemistry staining of dermal fibroblast for fibroblast marker (a. 1:200; b. 1:500), x10 objective magnification under phase contrast.

a. 1:200

b. 1:500



Figure 8: Immunocytochemistry staining of dermal fibroblast for Vimentin (1:400), x10 objective magnification (Staining intensity: +++ high).



Figure 9: Immunocytochemistry staining of dermal fibroblast for Melanocortin 1 receptor (1:10), x10 objective magnification (staining intensity ++ medium).

Discussion:

Fibroblast cells proliferate, migrate, synthesize ECM macromolecules, secrete growth factors, they are professionals in movement, ECM production, and deposition, and in a multitude of filopodial activities that enable them to shape the ECM and pull other cells. They make collagens, glycosaminoglycans, reticular and elastic fibres, and glycoproteins found in the extracellular matrix and tissue damage stimulates them and induces their proliferation. Stimulation of fibroblast migration by cytokines promotes their

migration and enhances the rate of wound healing (Schreier, et al, 1993). This study provides strong supporting evidence for an active involvement of the fibroblast cells, particularly DS cells in the wound healing. Using the transwell assay, human fibroblasts shows high rate of movement through the 8 µm pore membrane, this movement observed specially when combined with extracellular matrix protein (Fibronectin), but it is also significantly observed when combined with POMC

peptides. Cell migration can be regulated by peptides to ensure a fast formation of new connective tissue at the wound site. After the mechanical scratch of the monolayer, cells move forwards, coordinately as a sheet, to fill up the space without the requirement for cell division (Hall, 2005). The scratch assay shows successful results, revealed a significant high migration capacity of human fibroblast cells into the injured area. There was some variation in the cellular migration according to the media and the chemoattractant substance (POMC peptides). The enhanced healing or closure of the scratched area noted following the addition of POMC peptides, but the more potent chemotactic agent is the serum followed by the ACTH 10-7 M. It can be seen that in the first 72 hours from the scratch applied there is a rapid rate of cell migration, the cells become retracted at the wound edge, re-spread and extend filopodia into the scratch space, where they are in contact with their neighbours. Interestingly at 144 hours from the scratch wounding (the 7th day) there was significantly faster migration of the cells (leading to injury closure) i.e. as in monolayer feed with supplement medium. The fibroblasts express the intermediate filament protein vimentin which is the earliest to be expressed during cell differentiation, a feature used as a marker to distinguish their mesodermal origin. Vimentin is secreted in a large range of mesenchymal cell types

including fibroblasts, endothelial cells etc, and in a number of other cell types derived from mesoderm and mesoderm derived epithelia. Vimentin is found in various different tumours but is particularly expressed in those originated from mesenchymal cells. To investigate that it was important to know about the immunocytochemistry of the fibroblast cells in vitro, the fibroblast cells were tested positive for the protein vimentin. The POMC peptide products α -MSH and ACTH bind to specific receptors known as the melanocortin (MC) receptors. There is increasing evidence that the MC receptor type 1 (MC-1R) is expressed in vitro by fibroblasts as well as by several other cutaneous cell types besides melanocytes and keratinocytes (Böhme et al, 1999). The fibroblast cells in vitro were positive for MC1R. The MCR subtypes have been individually expressed in mammalian cell lines and their binding to MSH peptides has been characterized. Site-directed mutagenesis and molecular modelling of the MC1R has revealed a putative interaction of individual amino acids in the receptor with the MSH ligands. In conclusion, results presented in this work provide that the fibroblast subtypes DS play a key role in the wound healing specially when stimulated by POMC peptides. It was concluded also that the extracellular matrix protein (fibronectin) increased the migration rate of the fibroblasts.

Reference:

1. Böhm M, Metze D, Schulte U, Becher E, Luger TA, and Brzoska T. (1999): Detection of melanocortin-1 receptor antigenicity on human skin cells in culture and in situ. *Exp. Dermatol.* 8, 453-461
2. Böhm M and Luger TA. (2000): The Role of Melanocortins in Skin Homeostasis. *Horm. Res.* 54, 287-293
3. Branton, M. H. & Kopp, J. B. (1999) TGF-beta and fibrosis. *Microbes Infect. The American Journal of Surgery* 1:1349-1365.
4. Catania A and Lipton JM. (1993): α -Melanocyte-stimulating hormone in the modulation of host reactions. *Endocr Rev* 14: 564
5. Eichler MJ and Carlson MA (2006): Modeling dermal granulation tissue with the linear fibroblast-populated collagen matrix: a comparison with the round matrix model. *J Dermatol Sci.* 41(2): 97-108
6. Eves P, Layton C, Hedley S, Dawson RA, Wagner M, Morandini R, Ghanem G, Mac Neil S. (2000): Characterisation of an *in vitro* model of human melanoma invasion based on reconstructed human skin. *Br J Dermatol* 142: 210
7. Freinkel, R. K. Woodley, D. T. eds. (2001): *The Biology of the Skin*. Parthenon Publishing New York, NY 191-297
8. Grinnell, F. (1994): "Fibroblasts, myofibroblasts, and wound contraction". *J. Cell Biol* 124 (4): 401-404.
9. Hall A. (2005): Rho GTPases and the control of cell behaviour, MRC Laboratory for Molecular Cell Biology and Department of Biochemistry and Molecular Biology, Cancer Research UK Oncogene and Signal Transduction. *Biochemical Society Transactions* 33: 891-895
10. Hinz B (2006): The role of matrix adhesions in myofibroblast force perception and transmission. *European Journal of Cell Biology*, 85 (3-4): 175-181
11. Jahoda CA, Oliver RF, Reynolds AJ, et al. (2001b): Trans-species hair growth induction by human hair follicle dermal papillae. *Exp Dermatol.* 10: 229-237
12. Jahoda CAB, and Amanda J Reynolds. (2001a): Hair follicle dermal sheath cells: unsung participants in wound healing. *The Lancet*, 358 (9291), 1445-1448
13. Luger TA, Brzoska T, Scholzen TE, Kalden DH, Sunderkotter C, Armstrong C, Ansel J. (2000): The role of α -MSH as a modulator of cutaneous inflammation. *Ann. NY Acad. Sci.* 917: 232

14. Luger TA, Schwarz T. (1995): Effects of UV-light on cytokines and neuroendocrine hormones. In: Krutmann J, Elmers C (eds). *Photoimmunology* Oxford: Blackwell, 55-76
15. MacNeil S. (2007): Progress and opportunities for tissue-engineered skin. *Nature* 445 (7130): 874-80.
16. Millington G. (2006): Proopiomelanocortin (POMC): the cutaneous roles of its melanocortin products and receptors. *ClinExpDermatol* 31, 407-412.
17. Montagna W, Yun J, Formisano V (1964): Histology and cytochemistry of human skin. XXX. Cholinesterase-containing nerves in face. *Arch Dermatol* 90:526-529
18. Mulvaney M, Harrington A, James WD., Davies LB, Quick and Siegel (1994): *Cutaneous trauma and its treatment*. Military dermatology, Washington, DC: Office of the surgeon General at TMM publication. 143-156
19. Romo T, Presti PM, Yalamanchili HR. (2006): Medpor alternative for microtia repair. *Facial Plast Surg Clin North Am.* 14(2):129-36
20. Ruta M, Saleniece K, Riekstina U, Krigere L, Tirzitis G and Ancans J. (2007): Betulin binds to melanocortin receptors and antagonizes α -melanocyte stimulating hormone induced cAMP generation in mouse melanoma cells. *Cell Biochemistry and Function*, 25 (5): 591-596
21. Schreier T, Degen E, Baschong W. (1993): Fibroblast migration and proliferation during in vitro wound healing. *Res Exp Med*, 193(4): 195-205
22. Slominski A, Wortsman J, Szczesniewski A. (2000a) Liquid chromatography mass spectrometry detection of corticotropin-releasing hormone and proopiomelanocortin-derived peptides in human skin. *J Clin Endocrinol Metab* 85:3575-3581
23. Slominski A, Wortsman J, Luger T, Paus R, Solomon S (2000b): Corticotropin releasing hormone and proopiomelanocortin involvement in the cutaneous response to stress. *Physiol Rev* 80:979-1020
24. Wahl, SM. (1999): TGF- β in the evolution and resolution of inflammatory and immune processes. *Introduction, Microbes Infect* 1: 1247-1249