

<https://doi.org/10.54361/ljmr.12.2.13>

Serum Lipids and urate in Cigarette smokers

Altayeb Elazomi¹, Sara E. Ibrahim², Elrasheed I. Mohamed¹, Salah B. Bahroun⁴,
H. I. Fabelbum² and Omar F. Idres³

¹Faculty of Medical Technology, University of Zawia, Libya.

²Sharq Elniel College, School of Medical Laboratory Science.

³Faculty of Science and Technology, University of Alneelain, Khartoum, Sudan.

⁴National Medical Research Center, Zawia, Libya.

Abstract

Objective :The present study was undertaken to evaluate lipid profile and urate in apparently healthy cigarette smokers and compare it with apparently healthy non- smokers in the fasting state. Also to demonstrate the possible effect of smoking on serum lipid profile and urate concentration. **Methods:** The study was conducted on 50 healthy cigarettes smokers and compared with 50 healthy age and weight matched non-obese non-smokers who served as controls. Subjects in both groups were in the age range of 30-60 yrs., having no history of alcohol abuse, or diseases like diabetes mellitus, hypertension, hepatic impairment, renal disease, gout, hypouricemia, and obesity, and were neither on drugs like β -blockers, lipid lowering drugs, or thiazide diuretics. Clinical data were obtained from the history and record on questionnaire sheet. The clinical assessment was done by a medical doctor. **Results:** It was revealed that mean S.TC (275 ± 25 mg/dl), LDL-C (179 ± 44 mg/dl), and TG (116 ± 29 mg/dl), were significantly higher in smokers ($p < 0.001$) as compared to non-smokers, i.e., mean S.TC (139 ± 25 mg/dl), LDL-C (89 ± 25), TG (93 ± 25). mg/dl ($p < 0.001$) as compared to non-smokers. Mean serum HDL-C (27 ± 6 mg/dl) was significantly lower in chronic smokers ($p < 0.001$) as compared to non-smokers, i.e., 40.0 ± 8 mg/dl. Mean serum Urate (5.2 ± 1.0 mg/dl) was significantly lower in chronic smokers ($p < 0.001$) as compared to non-smokers, i.e., 6.1 ± 1.1 mg/dl. **Conclusion:** Smoking produces adverse effects on lipid profile and, therefore increasing the cardiovascular disease risk. On the other hand the significant low level of serum urate in smokers attributed in significant oxidative stress and hence cardiovascular disease. Cigarette smoke may be benefit in reducing hyperuricemia risk.

Keywords: S.TC : Serum Total cholesterol; LDL-C : Low density lipoprotein cholesterol; HDL-C : High density lipoprotein cholesterol; TG : Triglycerides; urate.

Introduction:

Worldwide, the effect of smoking are estimated to kill 3 million people per year. This contracts with 0.2 million in 1950 and projections for 2025 of 10 million. ¹ Epidemiological evidence suggest that

cigarette smoking is the a major risk factor for chronic obstructive pulmonary disease such as chronic bronchitis and emphysema, carcinogenesis and cardiovascular disease ^{1,2,3}. Various mechanisms leading to lipid

alteration by smoking are: (a) nicotine stimulates sympathetic adrenal system leading to increased secretion of catecholamines resulting in increased lipolysis and increased concentration of plasma free fatty acids (FFA) which further result in increased secretion of hepatic FFAs and hepatic triglycerides along with VLDL-C in the blood stream^{4,5}; (b) Fall in estrogen levels occurs due to smoking which further leads to decreased HDL – cholesterol³⁵; (c) Presence of hyperinsulinaemia in smokers leads to increased cholesterol, LDL-C, VLDL-C, and TG due to decreased activity of lipoprotein lipase^{6,7}; (d) Consumption of a diet rich in fat and cholesterol as well as a diet low in fibre and cereal content by smokers^{4,8}. Also recent evidence suggest that oxidants present in the gas phase of

Material and Methods

50 cigarette smokers and 50 normal healthy non smokers (control) were selected from the teaching staff of Sudan University for Science and Technology. None of them had past or present history of renal, gout, The smokers were age matched to controls (30 – 60) and they were regular cigarette smokers for the last 8 years, consuming more 10 cigarette per day. Clinical data were obtained from the history and record on questionnaire sheet. Clinical assessment was

Results

cigarette smoke are involved⁸. There is increasing experimental evidence that oxidation of low density lipoprotein cholesterol (LDL-C) plays a major role in the pathogenesis of coronary artery disease (CAD). However, results from clinical studies on LDL oxidation and CAD are not consistent¹⁰. Cigarette smokers have increased inflammatory responses that further enhance their oxidative stress.^{11,12} Since in humans, uric acid is the most abundant aqueous antioxidant, accounting for up to 60% of serum free radical scavenging capacity¹³ and is an important intracellular free radical scavenger during metabolic stress including smoking,^{14,15} therefore, measurement of its serum level reflects the antioxidant capacity.¹³

cardiac, hepatic, diabetes mellitus, hypertension endocrine disorders, obesity and on drugs like β -blockers, lipid lowering drugs, and thiazide diuretics.

done by a medical doctor. After overnight fasting, they were subjected to laboratory investigations including, lipid profile (serum cholesterol, triglycerides, HDL-C, LDL-C), and serum urate.

Table (1) and fig. (1) show, significant difference between the means of serum total cholesterol in the control group (n =50) and the study group (smokers). (n =50). (139 ± 25 versus 275 ± 51 mg/dl, $p < 0.001$). Fig (1) shows comparison of serum total cholesterol levels between control and smokers. Table (1) and fig. (2) show, significant difference between the means of serum LDL in the control group (n =50) and the study group (smokers). (n =50). (89 ± 23 versus 179 ± 44 mg/dl, $p < 0.001$). Fig (2) shows comparison of serum LDL levels between control and smokers.

Table (1) and fig. (3) show, significant difference between the means of serum triglyceride in the control group (n =50) and the study group (smokers). (n =50). (93 ± 25

versus 116 ± 29 mg/dl, $p < 0.001$). Fig (3) shows comparison of serum triglyceride levels between control and smokers.

Table (2) and fig. (4) show, significant difference between the means of serum HDL in the control group (n =50) and the study group (smokers). (n =50). (40 ± 6 versus 27 ± 8 mg/dl, $p < 0.001$). Fig (4) shows comparison of serum HDL levels between control and smokers.

Table (2) and fig. (5) show, significant difference between the means of serum urate in the control group (n =50) and the study group (smokers). (n =50). (6.1 ± 1.1 versus 5.2 ± 0.98 g/dl, $p < 0.001$). Fig (4) shows comparison of serum urate levels in control and smokers.

Variable	Control group n = 50	Study group n= 50	P
Serum cholesterol (mg/dl)	139 ± 25 (77 – 189)	275 ± 51 (177 – 415)	<0.001
Serum LDL	89 ± 23 (43 – 144)	179 ± 44 (82 – 273)	<0.001
Serum triglyceride	93 ± 25 (54 – 141)	116 ± 29 (58 – 186)	<0.001

Table (1) :Comparison of the means of serum total cholesterol, LDL, and triglyceride in the control group and the study group (smokers).

Variable	Control group n = 50	Study group n= 50	P
Serum HDL (mg/dl)	40 ± 6 (24 – 56)	27 ± 8 (11- 51)	<0.001

Serum urate (mg/dl)	6.1± 1 (4.4 – 10.6)	5.2± 0.98 (3.4 – 7.1)	<0.001
------------------------	---------------------	-----------------------	--------

Table (2): Comparison of the means of serum HDL, and urate in the control group and the study group (smokers).

- Table(1), and (2) show the mean ± S.D, range between brackets () and probability (p)

-t -test was used for comparison.




Figure (1): Comparison of levels of S. cholesterol (mg/dl) in control group and study group (smokers).




Figure (2): Comparison of levels of LDL- cholesterol (mg/dl) in control group and study group (smokers).




Figure (3): Comparison of levels of S. triglyceride (mg/dl) in control group and study group (smokers).




Figure (4): Comparison of levels of S. HDL- cholesterol (mg/dl) in control group and study group (smokers).




Figure (5): Comparison of levels of S. urate (mg/dl) in control group and study group (smokers).

Discussion

To our knowledge there appears to be no relevant published data on the effect of

smoking on blood chemistry among Sudanese, so comparison will be made with

studies made elsewhere. Prevalence of smoking varies considerably in different countries and we think it is increasing in Sudan. And this resembles the situation in other parts of Africa.^{16,17} Our study was done exclusively among the male teaching staff members of Sudan University for Science and Technology. In Sudan society female are reluctant to discuss the issue of smoking in an open manner, as smoking in young females is considered offending practice although it is accepted in old ones. It is revealed that triglycerides, LDL-C, TC were significantly higher in smokers as compared to non-smokers thereby revealing a direct dose response relationship^{8,18}. The mean serum total cholesterol in non-smokers (control) was 139.1 ± 25 mg/dl while it was significantly higher in smokers, i.e., 275 ± 15 mg/dl. These observations are in tune with the findings of other workers^{4,19,20}. Cigarette smoking substantially increases the risk of coronary heart disease and ischemic stroke^{22,23}. Serum LDL – cholesterol showed 101% increase, when the control group was compared to the study group (89 ± 23 versus 179 ± 44 mg/dl, $p < 0.001$). Serum LDL – cholesterol was highly significantly raised in the study group. This is in agreement with previous result published by many authors.^{2,9,16,24} The mean HDL-Cholesterol in non-smokers was $40. \pm 6$ and $27. \pm 8$ in smokers respectively (p value < 0.001). This finding is similar to

Conclusion

Is obvious from these results, cigarette smoking has a hyperlipidemic effect, which is reflected as raised serum levels of total cholesterol, LDL – cholesterol and triglyceride with reduced HDL – cholesterol,

that of Rosenson²² who reported that there is fall in HDL-C level by 3-5 mg/dl in smokers. The mean serum triglycerides levels in non-smokers and smokers were 93 ± 25 mg/dl and 116 ± 29 mg/dl, $p < 0.001$ respectively, serum triglyceride was highly significantly raised in the study group (smokers). However, triglyceride levels in study group (smokers) were found to be within the normal reference range.^{25,26} These findings suggest that smoking alters the lipid profile adversely causing dyslipidaemia in smokers and the changes become more marked with the number of cigarettes smoked²⁷ Smoking causes an increase in oxidized LDL-cholesterol level which plays the key role for atherosclerotic process¹⁹. A high level of LDL-C, VLDL-C and triglyceride are strongly associated with development of coronary artery disease while a low level of HDL-C remains a significant independent predictor of coronary artery disease^{29,30}. Serum urate on average, showed 14.8% reduction when the control group was compared to the study group (6.1 ± 1.1 versus 5.2 ± 1.0 mg/dl, $p < 0.001$), serum urate was highly reduced in the study group. This finding is in agreement with other studies that showed low serum uric acid in regular smokers^{30,32,34} and reduction of antioxidants including uric acid in smokers^{33,34} indicating that oxidative stress increases every time a cigarette is smoked.³¹

that increase the risk for atherosclerosis and ischemic heart disease. Serum levels of urate were found to be clearly reduced in cigarette smokers, this may attributed in significant oxidative stress and hence cardiovascular

disease. On other hand reduced serum urate gout.
may be benefit against hyperuricemia and

References

- 1- Bertteridge DJ, Morrel JM. Clinicians guide to lipids and Coronary Heart Disease. London : Chapman and Hall Medical 1998; 126 – 130.
- 2- Adlouni A, El-Messal M, Ghalim N, Saile R. Apolipoprotein and lipoprotein particles in Moroccan patients with previous myocardial infarction. *Int-Clin-Lab Res* 1997; 27 (4) :247 – 52.
- 3- Eiserich JP, Vander Vliet A, Hendelman GJ, Halliwell B, Cross CE. Dietary antioxidants and cigarette smoke induced bimolecular damage : a complex interaction. *Am J clin Nutr* 1995 Dec;62 (16) : 1490 – 1500.
- 4- Muscat JE, Harris RE et al. Cigarette smoking and plasma cholesterol. *Am Heart J* 1991; 121: 141-7.
- 5- Simons LA, Simons J, Jones AS. The interaction of body weight, age, cigarette smoking and hormone usage with blood pressure and plasma lipids in an Australian community. *Aus NZ J Med* 1984; 14: 215-21.
- 6- Stalder M, Pometta B, Suenram A. Relationship between plasma insulin levels and HDL-cholesterol in healthy men. *Diabetologia* 1984; 21: 544-8.
- 7- Reaven GM. Role of insulin resistance in human disease. *Diabetes*1988; 37: 595-607.
- 8- Wynder EL, Harris et al. Population screening for plasma cholesterol. Community based results from Connecticut. *Am Heart J* 1989; 117: 649-56.
- 9- Mc Call MR, Van Den Berg JJ, Kuypers FA, Tribble DL, Krauss RM, Knoff LJ, Forte TM. Modification of LACT activity and HDL structure; New Links between cigarette smoke and coronary heart disease risk. *Arterioscler thromb* 1994 Feb; 14 (2) : 248 – 53.
- 10- Halevy D, Thiery J, Nagel D, Arnold S, Erdmman E, Holfing B, Cremer P, Seidel D. Increased oxidation of HDL in patients with coronary artery disease is independent from dietary vitamin E and C. *Arterioscler- Throm- Vas- Biol.*1997 Jul 17 (7) :1432 – 7.
- 11- Alberg A. The influence of cigarette smoking on circulating concentrations of antioxidant micronutrients. *Toxicology* 2002. Nov;180(2):121-137.
- 12- Brown KM, Morrice PC, Duthie GG. Erythrocyte vitamin E and plasma ascorbate concentrations in relation to erythrocyte peroxidation in smokers and nonsmokers: dose response to vitamin E supplementation. *Am J Clin Nutr* 1997. Feb;65(2):496-502.

- 13- Maxwell SR, Thomason H, Sandler D, Leguen C, Baxter MA, Thorpe GH, et al. Antioxidant status in patients with uncomplicated insulin-dependent and non-insulin-dependent diabetes mellitus. *Eur J Clin Invest* 1997. Jun;27(6):484-490.
- 14- Mathru M, Dries DJ, Barnes L, Tonino P, Sukhani R, Rooney MW. Tourniquet-induced exsanguination in patients requiring lower limb surgery. An ischemia-reperfusion model of oxidant and antioxidant metabolism. *Anesthesiology* 1996. Jan;84(1):14-22.
- 15- Hellsten Y, Tullson PC, Richter EA, Bangsbo J. Oxidation of urate in human skeletal muscle during exercise. *Free Radic Biol Med* 1997;22(1-2):169-174.
- 16- Leonard L. Cigarette smoking and perception about smoking and health in chat. *E. Afr. Med. J* 1996; 73 : 509 – 512.
- 17- Nath UR. Smoking : Third world alert. London : Oxford University press 1996; 250 – 53.
- 18- Kannel WB. Update on the risk of cigarette smoking in coronary artery disease. *Am Heart J* 1981; 101: 319-28.
- 19- MJOS OD. Lipid effects of smoking. *Am Heart J* 1988; 115: 272-5
Rustogi R, Shrivastva SSI et al Lipid profile in smokers. *JAPI* 1989; 37 (12): 764-7
- 20- Austin MA. Plasma triglycerides and coronary heart disease. *Arterio Throm* 1991; 11: 2-14
- 21- Rosenson RS. Low level of HDL-cholesterol (Hypoalphalipoproteinemia). An approach to management. *Arch Intern Med* 1993; 153(13): 1528-40.
- 22- Berliner J, Navab M, Fogelman AM et al. Atherosclerosis: Basic mechanism-oxidation, inflammation and genetics. *Circulation* 1986; 5 (Supp): 5G-10G
- 23- Zimmerman M, Mc Geachie J. The effect of nicotine on aortic endothelial cell turnover. *Atherosclerosis* 1985; 58: 39-47.
- 24- The Expert Panel Report of the National Cholesterol Education Program. Expert panel on detection, evaluation and treatment of high blood cholesterol in adults. *Arch Intern Med* 1998; 148: 36-69.
- 25- Stampfer MJ, Sacks FM, Simonetta S et al. A prospective study of cholesterol apolipoproteins and the risk of myocardial infarction. *N Engl J Med* 1991; 325: 373-80.
- 26- Calver A, Collier J, Moncada S, Vallance P. Effect of local intra-arterial NG-monomethyl-L-arginine in patients with hypertension: the nitric oxide dilator mechanism appears abnormal. *J Hypertens* 1992. Sep;10(9):1025-1031.
- 27- Karnath B. Smoking cessation. *Am J Med* 2002. Apr;112(5):399-405.

- 28- Pryor WA. Cigarette smoke radicals and the role of free radicals in chemical carcinogenicity. *Environ Health Perspect* 1997. Jun;105(Suppl 4):875-882.
- 29- The Expert Panel Report of the National Cholesterol Education Program. Expert panel on detection, evaluation and treatment of high blood cholesterol in adults. *Arch Intern Med* 1998; 148: 36-69.
- 30- Tsuchiya M, Asada A, Kasahara E, Sato EF, Shindo M, Inoue M. Smoking a single cigarette rapidly reduces combined concentrations of nitrate and nitrite and concentrations of antioxidants in plasma. *Circulation* 2002. Mar;105(10):1155-1157.
- 31- Timimi FK, Ting HH, Haley EA, Roddy MA, Ganz P, Creager MA. Vitamin C improves endothelium-dependent vasodilation in patients with insulin-dependent diabetes mellitus. *J Am Coll Cardiol* 1998. Mar;31(3):552-557.
- 32- Dietrich M, Block G, Norkus EP, Hudes M, Traber MG, Cross CE, et al. Smoking and exposure to environmental tobacco smoke decrease some plasma antioxidants and increase gamma-tocopherol in vivo after adjustment for dietary antioxidant intakes. *Am J Clin Nutr* 2003. Jan;77(1):160-166.
- 33- Goraca A, Skibska B. Plasma antioxidant status in healthy smoking and non-smoking men. *Bratisl Lek Listy* 2005;106(10):301-306.
- 34- Nysoven EL. Ascorbate and urate smokers. *J clin. Nut.* 1995 Dec; 62 : 1490 – 1505.
- 35- Benowitz HL. Pharmacologic aspects of cigarette smoking and nicotine addiction. *New Engl J Med* 1998; 319: 318-30.